

## White Dwarfs from the SDSS: 90 Prime – Goin’ Deep in the White Dwarf Luminosity Function

James Liebert

*Steward Observatory, University of Arizona, Tucson, AZ 85721, USA*

Mukremin Kilic, Kurtis A. Williams, Ted von Hippel, and Don Winget

*Dept. Astronomy, University of Texas, 1 University Station, Austin TX 78712, USA*

Jeff Munn, Hugh C. Harris, and Stephen Levine

*U.S. Naval Observatory, Flagstaff Station, P.O. Box 1149, Flagstaff AZ 86002-1149, USA*

Travis S. Metcalfe

*High Altitude Observatory, NCAR, Mesa Laboratory, 1850 Table Mesa Dr., Boulder CO 80307-3000, USA*

**Abstract.** A large sample of cool white dwarfs from the Sloan Digital Sky Survey has been used to construct a luminosity function (LF), with the selection based on the method of reduced proper motion. While the sample is 60 times larger than that from our Luyten Half Second proper motion survey, its sensitivity to the region past the peak of the LF is severely limited by the necessity to match positions with detections from Palomar plates. A new survey to realize the full limiting magnitudes of the SDSS is described.

### 1. Introduction

A new disk white dwarf luminosity function (WDLF) has been constructed from Luyten Half Second (LHS) white dwarfs with motions having a minimum value of  $\mu > 0.6''/\text{yr}$  (Dahn, Harris, Leggett, & Liebert 2007, in preparation). While about doubling the size of the 1988 sample, it still includes only 92 cool white dwarfs. To eliminate spurious matches, the selection required measurement of reasonable astrometric positions from at least four Palomar Observatory Sky Survey first (POSS1) and second generation (POSS2) plates. The limiting magnitudes and corresponding search volumes were accordingly modest.

This LHS WDLF and a detail of the cool end are shown in Figure 1. The sharp peak before the plunge is real, and is believed to be caused by the “convective coupling” with the degenerate core, and a corresponding rapid release of interior energy (Fontaine, Brassard, & Bergeron 2001). Despite the small total number of stars, note that the last three half magnitude bins have 16, 10, and 3 stars, respectively. The plunge is therefore not measured with great statistical significance, and it has long been known that a sample at least several times larger has been needed (Wood & Oswalt 1998).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2007</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2007 to 00-00-2007</b>	
4. TITLE AND SUBTITLE <b>White Dwarfs From the SDSS: 90 Prime --Goin' Deep in the White Dwarf Luminosity Function</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>U.S. Naval Observatory,Flagstaff Station,P.O. Box 1149,Flagstaff,AZ,86002-1149</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			




Figure 1. Luminosity Function from 92 White Dwarfs in the Luyten Half Second Survey with  $\mu > 0.6''/\text{yr}$ , in 1/4 bolometric magnitude bins (filled circles). Hot stars (smaller triangles) are taken from the old study of DA stars from the Palomar Green Survey (Fleming, Liebert, & Green 1986).

## 2. The Sloan Digital Sky Survey

The SDSS has resulted in a dramatic several-fold increase in the known number of white dwarfs with spectral classifications. Over 9,300 are included in the Eisenstein et al. (2006) catalog. The great majority of these, however, are hot DAs and DBs. This is because hot stars are targeted for followup SDSS spectra, but cool white dwarfs lie too near to the locus of cool main sequence stars and are not targeted.

A new, much larger WDLF sample has been constructed by selecting stars based on reduced proper motion (RPM) –  $H_g = g + 5 \log \mu + 5$ . This is a proxy for absolute magnitude if the population sampled is kinematically similar. Specifically, the distribution of tangential velocities needs to be similar, since  $v_{\text{tan}} = 4.738 \mu d$ , where  $v_{\text{tan}}$  is in km/s and  $d$  is the distance in parsecs. Harris et al. (2006) published the WDLF and sample, based on reduced proper motion selection. To confirm where the boundaries between the bands of white dwarfs and main sequence subdwarfs lie, spectra obtained with the Hobby-Eberly and 6.5-m MMT were obtained for a subset published in Kilic et al. (2006).

In Figure 2 a RPM diagram is shown, using the SDSS  $g-i$  color. The dense contours to the right are a huge number of lower main sequence stars of the disk population. The main sequence stars of the local halo form a diagonal band off the lower left of the dense contours. Spectra confirming late-type subdwarf stars (sdK, sdM) are indicated with (red) squares. The white dwarfs form a broader diagonal band below and to the left of the subdwarfs. Those with spectral classifications – most hot ones from SDSS – are plotted as (blue) triangles. Black curves are the expected loci for white dwarfs with different mean  $v_{\text{tan}}$  values – the




Figure 2. A reduced proper motion diagram,  $H_g = g + 5 \log \mu + 5$  vs.  $g - i$  from the SDSS sample. The triangles identify spectroscopically-confirmed white dwarfs. Squares are late subdwarfs, or Population II main sequence stars. The broad contours are discussed in the text. The curves track the expected positions of DA stars with the labeled tangential velocity ( $V_T$ ).

higher the velocity, the higher the RPM at a given color. The color predictions are from the models kindly provided by P. Bergeron, assuming pure hydrogen atmospheric composition. As a result of the onset of the collision-induced  $H_2$  dipole absorption (CIA), the  $i$  band gets depressed for the coolest white dwarfs, and the curves are predicted to turn sharply blueward. Weaker CIA opacity effects have been measured in warmer white dwarfs in the near-infrared  $JHK$  bands (Bergeron, Ruiz, & Leggett 1997).

A few candidate cool CIA white dwarfs in the  $i$  band are identifiable near the bottom – middle of the figure. The limiting POSS magnitudes no doubt inhibit our ability to find many. Some were previously found in some SDSS-obtained spectra by Gates et al. (2004). When the signature is strong, the colors can be pulled away from the stellar locus, increasing the chance that the project would target them for spectra. The number of such objects, however, is small. By no means is the evidence from SDSS indicating that a large sample of such white dwarfs inhabit the Galactic halo, at least near the Sun.

A set of MMT spectra of confirmed cool white dwarfs from this RPM diagram, taken from Kilic et al. (2006), are shown in Figure 3. The vast majority of these stars do not show unusual spectra like the Gates et al. stars, or the strong-lined DZ and DQ white dwarfs illustrated in Harris et al. (2003). As can




Figure 3. 6.5-m MMT spectra of some cool white dwarfs confirmed in the sample

be seen, most are simple DC or DA stars. One star does show weak DZ metal features. Their spread in  $g - i$  and other SDSS colors is generally modest.

In Figure 4 the new WDLF constructed from this sample with 5,794 stars is shown. However, only two of these lie in the crucial last half-magnitude bin. Why is this? To measure a reduced proper motion, one needs to link the SDSS with other epochs to establish whether a consistent amplitude and position angle indicate a valid motion. We found that four matches were needed with this number of POSS plates. Thus, the selection of SDSS stars was also limited by the limits of the POSS. At the faintest absolute magnitudes, a sharp cut in the number of candidates thus occurred.




Figure 4. The luminosity function (LF) from the SDSS sample, in 1/2 magnitude bins, with the number of stars per bin labeled. Also shown are the new LF for hot DA white dwarfs from Liebert, Bergeron, & Holberg (2005), and the old LF of Leggett, Ruiz, & Bergeron (1998) from the 1988 LHS sample, again connected to the Fleming, Liebert, & Green (1986) sample of hot stars.

### 3. 90 Prime

In order to realize the limiting magnitudes of SDSS which extends about two magnitudes fainter than the POSS, it is obviously necessary to obtain at least one second epoch reaching similar magnitudes. For this purpose we have begun a large survey using a new prime focus imager on the Steward Observatory Bok 2.3-m (or 90-inch) reflector, described in Williams et al. (2004). The principal investigator is Ed Olszewski. The imager has an array of four thinned Lockheed  $4096 \times 4096$  pixel CCDs. The f/2.98 system provides a plate scale of  $0.45''/\text{pixel}$  and a total field-of-view of  $1.16^\circ \times 1.16^\circ$ .

We have estimated that several thousand square degrees of SDSS fields need to be repeated if we are to increase adequately the numbers of white dwarfs beyond the WDLF peak. This entails a huge observing project, estimated to be as many as 100 successful nights, assuming about 50 fields per night. An SDSS  $r$  band filter is available with the instrument, and only this band is being used. Early measurements indicate that valid proper motion stars may be found with only two epochs separated by several years in time. Both point spread functions are obtained with CCDs, so that more accurate astrometry is obtainable than with plates.

Eventually, additional epochs will be obtained by other projects. The most immediate is from the Pan-Starrs project (Hodapp et al. 2004). Using just the one telescope prototype to what is intended to be a 4-telescope system, an all-sky survey will be undertaken of the  $3\pi$  steradians of sky accessible from Maui, Hawaii. Five bands will be covered, including an “ $r$ ” band. Multiple measurements of the same sky fields are planned, resulting in the opportunity for multiple epochs for astrometry. Similar (deeper) multi-epoch imaging of astrometric quality may eventually be obtained when the Large Synoptic Survey Telescope (Tyson 2005) is built. However, the site selected for LSST is in Chile, affording less overlap with SDSS survey fields.

**Acknowledgments.** The authors grateful for additional support from the National Science Foundation through AST-0307321 and AST-0607480. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

## References

- Bergeron, P., Ruiz, M. T., & Leggett, S. K. 1997, *ApJS*, 108, 339  
Eisenstein, D. J., et al. 2006, *ApJS*, 167, 40  
Fleming, T. A., Liebert, J., & Green, R. F. 1986, *ApJ*, 308, 176  
Fontaine, G., Brassard, P., & Bergeron, P. 2001, *PASP*, 113, 409  
Gates, E., et al. 2004, *ApJ*, 612, 129  
Harris, H. C., Liebert, J., Kleinman, S. J., et al. 2003, *AJ*, 126, 1023  
Harris, H. C., Munn, J. C., Kilic, M., et al. 2006, *AJ*, 131, 571  
Hodapp, K. W., Kaiser, N., et al. 2004, *Astron. Nach.*, 325, 636  
Kilic, M., Munn, J., Harris, H. C., Liebert, J., von Hippel, T., Williams, K. A., Metcalfe, T. S., Winget, D. E., & Levine, S. E. 2006, *AJ*, 131, 582  
Leggett, S. K., Ruiz, M. T., & Bergeron, P. 1998, *ApJ*, 497, 294  
Tyson, A., 2005, in *ASP Conf. Ser.* 339, *Observing Dark Energy*, eds. S. C. Wolff & T. R. Lauer, (San Francisco: ASP), 95  
Williams, G. G., Olszewski, E., Lesser, M. P., & Burge, J. H. 2004, *SPIE*, 5492, 787  
Wood, M. A., & Oswalt, T. D. 1998, *ApJ*, 497, 870